

Ministry of Water Resources and Meteorology



Water Accounting Plus Framework: A Cambodia Case Study to Improve Water Resources Management


Prepared by : Hun Sothy, Mao Hak and Elga Salvadore


02-05 Oct 2018, Manila, Philippines

This is not an ADB material. The views expressed in this document are the views of the author/s and/or their organizations and do not necessarily reflect the views or policies of the Asian Development Bank, or its Board of Governors, or the governments they represent. ADB does not guarantee the accuracy and/or completeness of the material's contents, and accepts no responsibility for any direct or indirect consequence of their use or reliance, whether wholly or partially. Please feel free to contact the authors directly should you have queries.

Outline

- Introduction: River networks in Cambodia and Issues
- ADB/IHE Delft support: Water Accounting Plus (WA+) Framework
- Examples of WA+ results: performance of remote sensing data and models
- Conclusions and Next steps






Key Issues


- Water for agriculture is a key issue in Cambodia as crop production contributes to 54% of the sector GDP. Increase of rice exports are vital for Cambodia economy.
- Cambodia is highly dependent on water coming from upstream countries.
- The decrease in forest cover cause negative impact to the water retention
- Dry season: water scarcity, food security
- Wet season: flood, flash flood

Land Use and Land Cover for the Tonle Sap



UNESCO-IHE Institute for Water Education


Rainfall Remote Sensing





CONSUMPTIVE USE


Water withdrawn for a process that does not return back into the basin. It comprises of evapotranspiration (ET), water pollution and water incorporated into products.




During the wet season, the accumulated flow is higher (from 100 to 3000 m3/s at Lumphat) and more pixels contribute to the flow in the major streams

August

January 2006





Relatively precise results of modelled data

Basins significantly depend on upstream flow


Basin	Qsurf in / Net Inflow [%]
Tonle Sap	22.0
Three S	42.9
Coastal Catchments	0.0
Mekong Delta	89.8
Upper Mekong	92.0

Basins utilize less than 60% of the available water resource

Basin	Available [km3/yr]	Utilizable [km3/yr]	Utilizable/Available [%]
Tonle Sap	79.4	40.5	51.0
Three S	82.9	37.2	44.9
Coastal Catchments	30.3	9.6	31.7
Mekong Delta	526.6	274.9	52.2
Upper Mekong	472.0	279.9	59.3

Agricultural Services

- Values of land productivity (kg/ha) and water productivity (kg/m³)are generally low
- The 3S basin and Banteay meanchey province seems to contain the best performing food production area.



Related to Capacity Building

- Understanding concepts of WA⁺
- Hydrology, GIS and Remote Sensing
- Global hydrological modelling
- Obtain/process Remote sensing data
- Calculate/ Interpret P-ET
- Use remote sensing for WA⁺
- From WA⁺ sheets to diagnosis

Conclusion

- The power/accuracy of the data from satellite
- Improve the knowledge of technical staff on remote sensing product
- Understand the advantage of using remote sensing to complement the scarce ground measurements
- Help government and other key decision-maker on the use of this system for taking smart decisions.

Next Steps

- Incorporate WA+ framework in the National Water Resources Information System in the National Water Resources Data Center
- Continue capacity building for MOWRAM (and MAFF + MOE) in WA+
- Government technical staff to operate WA+ framework by 2025
- Government senior staff to use WA+ for planning, management, and monitoring of water resources
- Consider climate change in the WA+ framework for planning purpose?

Thank you very much for your attention