

# Regional Flyway Initiative - Site Study DRAFT

November 2025

# RFI Priority Site · Boeung Prek Lapouv (Sarus Crane Reserve)

Prepared by Ding Li Yong & Mike Crosby, with additional input from Bou Vorsak, Billy Fairburn, Kelvin Peh, Evelyn Pina-Covarrubias, Stefano Barchiesi and Radhika Bhargava BirdLife International, University of Southampton, National University of Singapore

Asian Development Bank

Disclaimer: The views expressed in this document are the views of the author/s and do not necessarily reflect the views or policies of the Asian Development Bank, or its Board of Governors, or the governments they represent. ADB does not guarantee the accuracy of the data included in this document and accepts no responsibility for any consequence of their use. By making any designation of or reference to a particular territory or geographic area in this document, the Asian Development Bank does not intend to make any judgments as to the legal or other status of any territory or area.

### **General Site Information**

Cambodia				
Boeng Prek Lapouv ID011				
Borey Chulsar and Koh Andeth district, Takeo Province				
10.72° N, 105.03° E	Area (has)	8,312 ha		
Sarus Crane (ssp. sharpii), Great Yellow-breasted Bunting	er Adjutant, Black-1	tailed Godwit,		
gallery forests (<1%), open war and rice fields (61%).	ter with aquatic pl	ants (13.7%),		
_				
Agricultural and fishing activities site.	s, Infrastructural de	evelopment near		
<ul><li>☑ Protected Area</li><li>☐ Ramsar</li><li>☐ others (</li></ul>	Site ☐ Flyway Ne	etwork Site )		
Boeng Prek Lapouv		,		
Ministry of Environment, Fisheries Administration, Provincial and district agricultural office, Takeo Provincial Government (including Provincial Department of Environment), Borey Chulsar and Koh				
No				
Site management, agriculture ar	nd small-scale fishe	ries		
Up to 10 years (but at least 5 ye	ars)			
Agriculture (rice) and fisheries				
		_		
⊠ No □ Yes				
16,450,000				
<ul><li>☑ Loan</li><li>☐ Private Sector</li><li>(to be identified)</li></ul>				
	☐ Public- Private	e Partnership		
	Boeng Prek Lapouv  Borey Chulsar and Koh Andeth of 10.72° N, 105.03° E  Sarus Crane (ssp. sharpii), Great Yellow-breasted Bunting  Seasonally inundated grassland gallery forests (<1%), open ware and rice fields (61%).  Global and local climate regulatore, flood hazard regulatore, and pollination  Agricultural and fishing activities site.  Protected Area Ramsar others ( Boeng Prek Lapouv  Ministry of Environment, Fisher district agricultural office, Taked Provincial Department of Environment Andeth district government  No  Site management, agriculture and Up to 10 years (but at least 5 years) Agriculture (rice) and fisheries  The wetlands sustain about 5, including at least 12 in Borey ( No Yes  Increase in tourism activities at ecosystems and potentially resulted to be identified)	Boeng Prek Lapouv  Borey Chulsar and Koh Andeth district, Takeo Provided Sarus Crane (ssp. sharpii), Greater Adjutant, Black-Yellow-breasted Bunting  Seasonally inundated grasslands (24% of total argallery forests (<1%), open water with aquatic pland rice fields (61%).  Global and local climate regulation. Fisheries, frod, rice, flood hazard regulation, water purificative treatment, and pollination  Agricultural and fishing activities, Infrastructural desite.  Protected Area Ramsar Site Flyway Notative (Boeng Prek Lapouv  Ministry of Environment, Fisheries Administration, district agricultural office, Takeo Provincial Governous Provincial Department of Environment), Borey Chulandeth district government  No  Site management, agriculture and small-scale fished Up to 10 years (but at least 5 years)  Agriculture (rice) and fisheries  The wetlands sustain about 5,000 households are including at least 12 in Borey Chulsar and Koh Are No Yes  Increase in tourism activities at the site may impact ecosystems and potentially result in disturbances to 16,450,000  Diagram Private Sector Public- Private		

### **Table of Contents**

General Site Information	3
Abbreviations	5
Executive summary	6
1. Background of the Regional Flyway Initiative	7
2. Site profile of Boeung Prek Lapouv (Sarus Crane Reserve)	8
3. Biodiversity value of Boeung Prek Lapouv (Sarus Crane Reserve)	10
3.1 Importance for migratory waterbird species	10
3.2. Other notable biodiversity	10
4. Ecosystem services	11
4.1. Ecosystem services provided by Boeung Prek Lapouv (Sarus Crane Re	eserve)11
4.2. Global climate regulating services	12
4.3. Flood mitigating services	12
5. Drivers of change and their potential impacts on Boeung Prek Lap Reserve)	•
5.1. Current drivers of change and their level of impact	13
5.2. Potential alternative state of Boeung Prek Lapouv (Sarus Crane Res drivers of change	•
5.3. Expected changes in the ecosystem services of Boeung Prek Lapouv (Sa	•
6. Capacity needs in Boeung Prek Lapouv (Sarus Crane Reserve)	16
7. Opportunities for RFI interventions	18
References	26
Annex 1. Supplementary information on flood mitigation services	28

#### **Abbreviations**

ADB Asian Development Bank AWC Asian Waterbird Census

BPL Bueng Prek Lpov

CSR Conservation Status Review

DoFWC Department of Freshwater Wetlands Conservation

DoF Department of Fisheries
DMC Developing Member Country

EAAFP East Asian-Australasian Flyway Partnership

FiA Fisheries Administration

GDANCP General Directorate of Administration for Nature Conservation and Protection

IBA Important Bird and Biodiversity Area

IUCN International Union for the Conservation of Nature

LCG Local conservation group

MAFF Ministry of Agriculture, Forestry and Fisheries

MOE Ministry of Environment NLC NatureLife Cambodia

NGO Non-governmental Organisation

PDoE Provincial Department of Environment

RFI Regional Flyway Initiative

TPA Takeo Provincial Administration

TESSA Toolkit for Ecosystem Services Assessment

USD United States Dollars

WCS Wildlife Conservation Society
WWT Wetlands and Wildfowl Trust

#### **Executive summary**

Cambodia's middle and lower floodplains along the Mekong River are home to some of very few untouched floodplain wetlands remaining in continental Southeast Asia. Spanning an area of nearly 8,000 ha, Bueng Prek Lapouv host important areas of seasonally flooded grassland and flooded forests and supports globally significant congregations of both migratory and resident waterbirds, the best known being the Sarus Crane, of which more than 10% of the regional population uses the site during the dry season. In recent years, surveys have also documented regular congregations of Greater Adjutant (NT), Spot-billed Pelican (NT) and very large congregations of the Black-tailed Godwit (NT) and Yellow-breasted Bunting (CR). BPL is designated as a Protected Landscape with the Ministry of Environment as its management authority. There are at least 12 villages across the two districts, Borey Chulsar and Koh Andeth that BPL falls under. Thousands of people rely on BPL for their livelihoods; an ecosystem services survey in 2012 found that almost 3,000 households (68% of the total) use the site for resources such as fish, edible plants, firewood, and for farming land, demonstrating the importance of the site for key provisioning services and regulating services from flood hazard control.

Local communities in and around BPL are dependent on small-scale fishing and dry-season rice farming as a major source of livelihood, together with nature-based tourism activities from mostly domestic visitors. Rice is also cultivated in the buffer zone, and experimental zones of BPL although the core conservation area (<2,200 ha) is relatively undisturbed. Major threats faced by BPL are land encroachment, illegal fishing, illegal (bird) trapping, pollution, and the overuse of water for growing rice during the dry season. Invasive plants such as the giant mimosa affect the site's wetland ecosystem but changing the plant composition within the wetland. Water pollution, especially from household waste and agro chemicals used in rice cultivation, may also have long-term impact on BPL's important fisheries.

To strengthen management and conservation of BPL there is a clear need for: improving legal protection and the overall management of the wetlands. This includes better zonation of key sites for livelihood and conservation activities, while addressing dry-season fires and invasive species. Although nature-based tourism activities at BPL at best limited, studies show that this is anticipated to grow over time, which will increase pressures on the site. This needs to be supported by a commensurate investment into sustainable nature-based tourism, including improvement of ecotourism infrastructure at BPL including signages, jetties and viewing structures. This then needs to be supported by targeted work to build the capacity of local communities and tourism businesses (e.g. skills-based training, hospitality training) to scale up tourism operations in an environmentally friendly manner. There are also major opportunities for rice agriculture transition with a focus on biodiversity-friendly rice farming approaches that has been piloted elsewhere in Cambodia, most famously through the 'Crane Rice' initiative in Kampot and 'Ibis Rice' at other inland sites.

#### 1. Background of the Regional Flyway Initiative

In July 2021, the Asian Development Bank made a commitment to develop a long-term Regional Flyway Initiative (RFI) in the East-Asian Australasian Flyway (EAAF) (Sovereign Project 55056-001) to protect and restore priority wetland ecosystems and the associated ecosystem services they provide in the EAAF, the most threatened migratory bird flyway globally. The Initiative is slated for implementation in nine ADB developing member countries (DMCs) in East, South and Southeast Asia: Mongolia, People's Republic of China (PRC), Bangladesh, Viet Nam, Cambodia, Philippines, Thailand, Malaysia and Indonesia. In 2023, the geographic scope of the RFI was further extended to two DMCs in Southeast Asia and the Pacific respectively, Lao PDR and Papua New Guinea.

The primary aim of the RFI is to enhance and expand the existing efforts in conserving and managing wetlands of the highest priority for migratory birds within the EAAF through innovative loan and grant financing, and at scale. Consultations and analyses over the development period help identify key interventions to strengthen the management of wetlands, enabling the implementation of nature-based solutions while strengthening biodiversity protection. Over time, the RFI seeks to leverage collaborative opportunities by developing partnerships among important stakeholders including national governments, civil society organizations, communities, regional organizations like the East Asian-Australasian Flyway Partnership (EAAFP), development agencies, the private sector, and other relevant entities.

Through the RFI Technical Assistance (TA) implemented over the RFI's development phase from 2021 to 2024, BirdLife International takes the lead in providing and coordinating technical support for development of the RFI. This is carried out in collaboration with the EAAFP and a consortium of international non-governmental organizations including Wetlands International and the Paulson Institute, as well as two universities, namely the University of Southampton, UK and the National University of Singapore. Over the development phase, the TA team undertook a site selection analysis to identify priority wetland sites in all 10 countries based on recent bird data benchmarked against internationally accepted criteria under the Convention on Wetlands of International Importance (or Ramsar Convention), EAAFP Flyway Network Sites and Important Bird and Biodiversity Areas (IBAs). The team further developed ecosystem services profiles for prioritised wetlands using a multi-pronged approach used the TESSA ecosystem services assessment tool, and data-driven modelling of water-based ecosystem services and stored carbon.

In Cambodia, a total of 15 wetland sites, including several Asian Waterbird Census (AWC) count sites, were initially assessed through data analysis and expert consultation, of which 12 were short-listed for assessment based on the available (recent) data. Of these pool of sites, nine (9) were defined and identified to be RFI priority sites on the basis that they support more than 1% the flyway population of at least one EAAF migratory waterbird species. Eight (8) of the RFI sites identified are inland wetlands, most notably a cluster of sites around the Tonle Sap Great Lake, such as Prek Toal, Ang Tropeang Thmar and Boeng Tonle Chmmar. A single coastal site was identified, i.e. Koh Kapik Ramsar Site where there have been extensive surveys of its biodiversity

to date, including surveys led by NatureLife Cambodia. 11 EAAF species exceeded the 1% threshold at the site level in Cambodia, with species such as the Masked Finfoot, Greater Adjutant and Sarus Crane (ssp. *sharpii*) at their highest congregations in Southeast Asia. Other species with important populations and/or congregations in Cambodia includes Spotted Greenshank, Black-headed Ibis and Painted Stork.

#### 2. Site profile of Boeung Prek Lapouv (Sarus Crane Reserve)

Location: Boeung Prek Lapouv is part of one of the largest areas of contiguous natural wetland (e.g. seasonally flooded grassland and mosaics of *Barringtonia*-dominated swamp forest) remaining in the Mekong Delta. The site is located in Koh Andeth and Borey Chulsar districts of Takeo Province in the western floodplain of the Bassac River, a distributary of the Mekong River. It borders Viet Nam to the south and east, where the natural habitat has been almost entirely converted to rice agriculture.

Area: Boeung Prek Lapouv (RFI site) covers an area of 8,312 ha

Altitude: 4-5 metres asl.

Geographical coordinates: 10.72° N, 105.03° E


Description of site: The Boeung Prek Lapouv (BPL) Protected Landscape is one of few remaining areas of seasonally inundated floodplain grasslands in the Cambodian part of the Lower Mekong Delta. The site is inundated for 3-4 months each year during the monsoon season, during which time a mat of floating vegetation forms on the surface of the water. Sophanna et al. (2019) identified four habitat types within BPL; they include seasonally inundated grasslands (24% of total area), shrubs and gallery forests (<1%), open water with aquatic plants (13.7%), and rice fields (61%). Grassland habitats throughout the region are under threat from agricultural conversion, with their hydrology significantly altered to facilitate rice growing, which poses a risk to flagship species including Sarus Crane Antigone antigone and Painted Stork Mycteria leucocephala. This wetlands of Boeng Prek Lapouv provide critical ecosystem services that is estimated to support the livelihoods of 12,000 people from 19 villages.

Site administration, management and land tenure: Boeung Prek Lapouv was established as a Conservation Area for Sarus Cranes in 2007, authorised under the Ministry of Agriculture, Forestry and Fisheries (MAFF), with a core conservation zone that included two community fisheries. The protected area was officially transferred to the Ministry of Environment (MoE) and re-designated as Protected Landscape in May 2016. Despite being established as a conservation area since 2007, Boeung Prek Lapouv has yet to be clearly demarcated, leading to uncertainty around land tenure and zonation and continuing encroachment. A governance structure has been developed for the protected area: within MoE, the institution responsible for taking the lead management role is the Department of Freshwater Wetlands Conservation (DoFWC) of the General Directorate of Administration for Nature Conservation and Protection (GDANCP) and the Provincial Department of Environment (PDoE). These organisations cooperate with Takeo provincial administration (TPA) and the District Administrative Authority of Takeo Province, and

WWT and BirdLife are the international NGO partners. A management plan for Boeung Prek Lapouv from 2014-2018 was published by WWT (2014). According to interviews with commune chiefs, most land in Boeung Prek Lapouv is presumed to be owned by local people, although few hold official land titles.

Social and economic values: About 22 villages (~5,000 households) use the wetlands at Boeung Prek Lapouv for rice farming and the collection of natural resources including fish, edible plants, firewood and grass. Sophanna (2017) found that most people there (97%) are involved in the cultivation of rice, raising cattle for sale (56%) and fishing (51%). There are tensions between the local population and transient Vietnamese farmers and fishers who cross the border to use the land. The widespread conversion of wetlands for rice cultivation is likely to result in a reduction in ecosystem values to local people because of a reduction in the availability of natural resources.

**Figure 1**. Map of the Boeng Prek Lpov Protected Landscape, showing its location in Cambodia (Map: NatureLife Cambodia)



#### 3. Biodiversity value of Boeung Prek Lapouv (Sarus Crane Reserve)

#### 3.1 Importance for migratory waterbird species

During the dry season, from December until May, Boeung Prek Lapouv supports an important non-breeding population of Sarus Crane *Grus antigone*, although their numbers there (and at other non-breeding sites for the species in Cambodia) have declined in recent years. Boeung Prek Lapouv also supports small numbers of several other globally threatened and near-threatened waterbird species, including Black-headed ibis *Threskiornis melanocephalus*, Greater adjutant *Leptoptilos dubius* (NT) and Lesser adjutant *Leptoptilos javanicus* (NT).

Count data from the 2017, 2019 and 2022 Asian Waterbird Census (AWC) was used in the RFI analysis for Boeung Prek Lapouv. The results of these counts were averaged for these three years, and then compared to the Conservation Status Review (CSR1) 1% population estimates to calculate a score for each species. The two migratory waterbird species were found to regularly exceed the 1% population estimates during these three years (Table 1), and the CSR1 scores for these species were summed to produce the overall site score.

**Table 1**. List of migratory species (based on the EAAFP list of species) with globally significant congregations in Boeng Prek Ppov wetlands.

Species name	IUCN	Average count	CSR1	CSR1 score
Sarus Crane Antigone antigone sharpii	VU	58.7	2	29.3
Painted Stork  Mycteria leucocephala	LC	73.5	70	1.05


#### 3.2. Other notable biodiversity

In addition to the waterbirds, Boeung Prek Lapouv supports small numbers of the globally threatened Bengal Florican *Houbaropsis bengalensis* (CR) which successful bred there in 2015 and there remains a small population as of 2025 (Vorsak, B. in litt.). The site is particularly notable for the large non-breeding population of the Yellow-breasted Bunting *Emberiza aureola* (CR), potentially the largest roost of the species in Cambodia, with more than 5,000 individuals counted there in 2022 (Ly et al. 2022).

#### 4. Ecosystem services

#### 4.1. Ecosystem services provided by Boeung Prek Lapouv (Sarus Crane Reserve)

The Boeung Prek Lapouv contains diverse wetland habitats, providing essential provisioning, regulating and cultural ecosystem services (Figure 2). The results from the RFI workshop highlights the top ecosystem services provided by the site, emphasising their essential and non-substitutable nature (Table 2). Provisioning services, particularly fresh water and food, benefit communities both within, adjacent to, and distant from the site. Regulating services, such as flood hazard regulation, water purification and waste treatment, and pollination, play a crucial role. Flood hazard regulation benefits communities within and adjacent to the site whereas water purification benefit communities within the site only. Pollination benefits communities at all distances.



**Figure 2**. List of ecosystem services provided by Boeung Prek Lapouv (Sarus Crane Reserve), as identified through stakeholder consultation at the Regional Flyway Initiative workshop.

**Table 2**. List of top ecosystem services provided by Boeung Prek Lapouv (Sarus Crane Reserve).

Ecosystem services	Essential	Benefits	Benefits to communities			Benefits to communities		Change	
	or non- substituta ble	Within the site	Adjacent to the site	Distant to the site	Past	Future			
Provisioning service	es								
Fresh water	Yes	<b>√</b>	✓	✓	Increase	Decrease			
Food	Yes	<b>√</b>	✓	<b>√</b>	Decrease	Increase			
Regulating services	5	ı		1		1			
Flood hazard regulation	Yes	✓	✓		Increase	Increase			
Water purification and waste treatment	Yes	✓			No change	No change			
Pollination	Yes	<b>√</b>	✓	<b>√</b>	Increase	No change			

#### 4.2. Global climate regulating services

Based on the look-up values from a FAO report (Dondini et al. 2023) and IPCC (2006), the amount of carbon stored in Boeung Prek Lapouv is estimated to range from 61,500 to 65,600 tonnes, while the annual carbon sequestration rate is estimated at 1,130 tonnes per year.

#### 4.3. Flood mitigating services

The flood mitigation services provided by Boeung Prek Lapouv were assessed using biophysical values only (see Table A1 and Annex 1 for details). When compared to both the average of the eight RFI inland sites and the average of all other inland wetlands in Cambodia (Table A2 in Annex 1), Boeung Prek Lapouv shows some consistent results in terms of benefits and beneficiaries:

- (1) for the average green storage capacity per sq. km of wetland, Boeung Prek Lapouv is above average (448 Giga Litres or GL of water per km² vs. 384 GL/km² for RFI inland sites and 399 GL/km² for all other inland wetlands);
- (2) for the average population uniquely benefitting from influential green storage upstream per sq. km of wetland, Boeung Prek Lapouv is also above average (139 vs. 81 people/km² RFI inland sites and 109 people/km² for all other inland wetlands); and

(3) for the average built-up area uniquely benefitting from influential green storage upstream per sq. km of wetland, Boeung Prek Lapouv is again well above average (8.40 ha/km² vs. 4.89 ha/km² for RFI inland sites and 5.24 ha/km² for all other inland wetlands).

## 5. Drivers of change and their potential impacts on Boeung Prek Lapouv (Sarus Crane Reserve)

#### 5.1. Current drivers of change and their level of impact

Stakeholders identified 31 drivers of change impacting Boeung Prek Lapouv, and their corresponding levels of impact on the wetland site (Table 3). High-impact drivers include the collection of terrestrial plants (non-timber), which significantly affects the natural vegetation and biodiversity. Drought conditions and prolonged droughts also pose substantial threats to the wetland's ecological balance, reducing water availability and affecting habitat quality. Fishing, killing, and harvesting of aquatic resources further exacerbate the impact on the site's biodiversity.


Medium-impact drivers consist of agricultural and forestry effluents, which degrade water quality and impact aquatic life. Annual and perennial non-timber crop production, dams and hydrological modifications, and habitat clearing contribute to changes in land use and habitat loss. Other medium-impact factors include invasive plant species, livestock farming, recreational activities and tourism, and research and education activities. Restoration for conservation has moderately modified the site, while water extraction, shipping lanes, canals, and air-borne pollutants add to the cumulative stress on the wetland's ecosystem.

Driver of change	Impact
Collecting terrestrial plants or plant products (non-timber)	
Drought conditions	⊔iah
Droughts	High
Fishing, killing and harvesting of aquatic resources	
Agricultural and forestry effluents	
Annual and perennial non-timber crop production	
Dams within or upstream of the wetland site, which alter the hydrological regime	
Dams, hydrological modification and water management/use	
Habitat clearing	
Invasive plant species	Medium
Livestock farming and grazing	Medium
Recreational activities and tourism	
Research, education and other work-related activities	
Restoration for conservation	
Shipping lanes and canals	
Water extraction/diversion within the wetland site or catchment	
Air-borne pollutants	
Fire and fire suppression	
Garbage and solid waste	
Habitat shifting and alteration	
Household sewage and urban wastewater from outside the wetland site	
Housing and settlement	
Hunting, killing and collecting of terrestrial animals	
Invasive animal species	Low
Loss of keystone species	
Pathogens	
Sewage and wastewater from wetland site facilities	
Storm and flooding	
Temperature extremes	
Vandalism, destructive activities or threats to staff and visitors	
Wood pulp and plantations	

**Table 3.** Drivers of change and their potential impact on the integrity of Boeung Prek Lapouv (Sarus Crane Reserve) based on consultations with stakeholders.


## 5.2. Potential alternative state of Boeung Prek Lapouv (Sarus Crane Reserve) under current drivers of change

Stakeholders defined the most plausible future alternative state (to 2035), and how this will translate to a net change in the cover of different types of wetland habitat types within this site (current habitat cover vs future alternative cover; Figure 3). The alternative state of the site assumes there will be no changes in the current drivers of change impacting the site, and the current management regime.



**Figure 3.** The proportional change in the extent of different habitat types between the current and alternative states of Boeung Prek Lapouv (Sarus Crane Reserve).

**Figure 4**. Major land cover types in BPL wetlands based on remotely sensed data (Map: Radhika Bhargava)



## 5.3. Expected changes in the ecosystem services of Boeung Prek Lapouv (Sarus Crane Reserve)

Stakeholders documented the future trends in the provision of ecosystem services in Boeung Prek Lapouv, indicating if the ecosystem services provided by this site (to 2035) will increase, decrease, or will remain stable if the current drivers of change impacting this site will continue in their present condition, with the intervention remains unchanged.

Figure 2 and Table 2 highlight that provisioning service for fresh water has seen an increase in the past, but a decrease is anticipated in the future. Cultivated good provision has increased in the past and this trend will continue in the future. Wild food provision has experienced a decline previously but is expected to increase moving forward. Flood hazard regulation has increased in the past, with further increases expected. Water purification has remained stable, showing no change in both the past and future. Pollination has increased in the past but is projected to remain unchanged in the future.

In the alternative state, the 40% gain wetland will result in a gain of stored carbon, estimated to be between 3,600 and 5,210 tonnes, and an increase in carbon sequestration rate (carbon accumulation) by approximately 253 tonnes per year.

A gain of 200.2 ha of freshwater, tree-dominated wetlands as presented in Table A5 is equivalent to about 2.4% of the total land use for the site, which is expected to result in roughly a 2.8% or 12 Giga Litre increase in green storage capacity per km² of wetland. This may amount to over 2 people and 0.11 hectares of built-up areas gaining flood mitigation benefits from Boeung Prek Lapouv per km² of wetland.

#### 6. Capacity needs in Boeung Prek Lapouv (Sarus Crane Reserve)

**Table 4.** Capacity needs for key stakeholders involved in the management of the BPL wetlands.

Stakeholder Group	Current role in the wetland management	Current capacity for sustainable wetland management	Capacity Development support needed to improve wetland management	Form of capacity development
Ministry of Environment	Oversee site management and	N/A	N/A	N/A
Provincial Department of Environment	policymaking.	N/A	N/A	N/A

Site management and rangers  Local authorities	Management of site, undertake patrols and enforcement. Engage local communities  Support with enforcement against illegal activities. Support site management	<ul> <li>GPS utilization</li> <li>Documentation of illegal cases</li> <li>Proficiency in law enforcement procedures</li> <li>Report writing skills</li> <li>Boat driving expertise</li> <li>Laptop operation</li> <li>Bird photography and identification</li> <li>Tour guiding capacity</li> <li>Awareness-raising</li> <li>Coordination skills</li> <li>Intervention in law enforcement</li> <li>Administrative proficiency</li> <li>Action plan development</li> </ul>	<ul> <li>GPS and GIS approaches</li> <li>SMART Patrol approaches</li> <li>Illegal case documentation</li> <li>Biodiversity monitoring</li> <li>First Aid</li> <li>Tourism management</li> <li>Site management</li> <li>Law enforcement</li> <li>Guidelines for wetland management</li> <li>Strengthen grievance mechanism</li> <li>Law enforcement</li> </ul>	<ul> <li>Training (online + In person)</li> <li>Site visit (local and oversea)</li> <li>Internship</li> <li>Equipment and tools (laptop, camera, GPS, Drone)</li> <li>Training</li> <li>Site visit</li> <li>Internship</li> </ul>
Other relevant ministries	N/A	N/A	N/A	N/A
Local communities	Direct beneficiaries of site	N/A	Support site management activities, enforcement and biodiversity monitoring.	<ul><li>Training programmes</li><li>Site visit</li></ul>

#### 7. Opportunities for RFI interventions

Boeung Prek Lapouv is one of very few large, floodplain wetland on the lower Mekong of Cambodia and supports globally significant congregations of both migratory and resident waterbirds, the best known being the Sarus Crane. BPL is designated as a Protected Landscape with the Ministry of Environment as its management authority. There are at least 12 villages across two districts at the site, and local communities are dependant on small-scale fishing and dry-season rice farming as a major source of livelihood, together with nature-based tourism activities from mostly domestic visitors. Rice is also cultivated in the buffer zone, and experimental zones of BPL although the core conservation area (<2,200 ha) is relatively undisturbed. Encroachment activities such as illegal land clearance (and land grab activities) have damaged or resulted in the loss of a significant part of BPL while fishing and agricultural activities is expected to contribute to water and solid waste pollution, which in turn may have long-term impact on BPL's important fisheries. To strengthen management and conservation of BPL there is a clear need for: (1) strengthening legal protection and management of the wetlands, including better zonation of key sites for livelihood and conservation activities, while addressing dry-season fires and invasive species such as giant mimosa which is now established across much of the region.

Although nature-based tourism activities at BPL is limited at present, studies show that this is anticipated to grow over time, which will benefit local communities but also increase pressures on the site. This needs to be supported by a commensurate investment into tourism, including (1) Improvement of ecotourism infrastructure at BPL including signages, jetties and viewing structures. This then needs to be supported by targeted work to build the capacity of local communities and tourism businesses (e.g. skills-based training, hospitality training) to scale up tourism operations in an environmentally friendly manner. There are also major opportunities for rice agriculture transition with a focus on biodiversity-friendly rice farming using approaches that has been trialled in Anlung Pring, Kampot. Existing consultations with local people has identified 'Crane Rice' as a potential intervention, in contributing to improved site management, while addressing broader livelihood issues for several local communities.

**Table 5**. List of proposed interventions for Bueng Prek Lpov wetlands and possible project indicators.

Intervention	Output	Cost (USD)	Timeframe	Indicators	Potential Stakeholder				
Component 1. Strengthening site management and protection of BPL.									
Scoping study of existing interventions and threats and disturbance at BPL, including targeted work on long-term threats (from climate change,	Assessment and scoping guidance to provide basis for proposed RFI interventions.	50,000	1 year	Assessment report with key threats identified and recommendations for improved management published, and	MOE  BPL management board  Consultancy companies				

encroachment and invasive species management).				disseminated to key stakeholders.	Conservation organisations ADB
Strengthen and expand site management and BPL protected landscape, including designation of a buffer.	Site management plan over short to medium- term for BPL developed, in consultation with key stakeholders  Establishment of a co- management framework, with stronger involvement of local stakeholders.  Measures to address and mitigate fire risk during the dry season defined and actively implemented.	100,000	2 years	Site management plan and zonation plan agreed by key stakeholders and endorsed by MOE  Stakeholder consultation meetings organised.  Meetings and workshops to engage local stakeholders for participatory processes, and national stakeholders for mandates on zonation.  Increased enforcement patrols against baselines, focusing on illegal bird hunting and land clearance.	MOE  BPL management board  Takeo provincial government  Borey Chulsar and Koh Andeth district government  Conservation organisations  Community groups (including LCGs)
Improve water management infrastructure to ensure water supply to BPL during the dry season to minimise the risk of fires.	Improve and expand water management infrastructure (including canals and ditches).	1,500,000	2-3 years	Drainage canals for water management repaired and maintained (including clearance of vegetation).  Localised dredging works to address siltation and vegetation.	MAFF BPL management board
Assess the extent of water, plastic and solid waste pollution in BPL (e.g. agro-chemicals from rice	Assessment study to determine the main causes of water pollution and potential approaches to address water pollution	100,000	2 years	Comprehensive study published and circulated to key stakeholders on water and solid waste pollution.	MOE  BPL management board  Conservation and academic

				organisations (including bird
				and community- focused NGOs
				Research institutions
Mimosa at BPL effectively controlled and no longer affecting sections of the BPL Core Zone identified to be important to biodiversity.	100,000	5 years	invasive species including giant mimosa effectively controlled in at least 20 ha of the wetlands.  At least one experimental plot for giant mimosa removal established.	BPL management board  Conservation and academic organisations (including bird and community- focused NGOs
Trials for managing giant mimosa established.	d monitoring	with a focus o	Invasive species monitoring and management plan developed and actively implemented by management board.  Management plan contribute to a part of the overall site management and zonation plan.	Research institutions  Community groups
	_		,	
	100,000	3 years	·	MOE
delineated for different usage.			approach strengthened and scaled up significantly against	BPL management board
Establishment			baselines.	Borey Chulsar
of a co- management framework, with stronger involvement of local stakeholders.			Co-management framework with representative from villages and conservation organisations.	and Koh Andeth district government  Conservation and academic organisations (including bird
	BPL effectively controlled and no longer affecting sections of the BPL Core Zone identified to be important to biodiversity.  Trials for managing giant mimosa established.   Zones in BPL area precisely delineated for different usage.  Establishment of a comanagement framework, with stronger involvement of local	Mimosa at BPL effectively controlled and no longer affecting sections of the BPL Core Zone identified to be important to biodiversity.  Trials for managing giant mimosa established.  Zones in BPL area precisely delineated for different usage.  Establishment of a co- management framework, with stronger involvement of local	Mimosa at BPL effectively controlled and no longer affecting sections of the BPL Core Zone identified to be important to biodiversity.  Trials for managing giant mimosa established.  Zones in BPL area precisely delineated for different usage.  Establishment of a co- management framework, with stronger involvement of local	Mimosa at BPL effectively controlled and no longer affecting sections of the BPL Core Zone identified to be important to biodiversity.  Trials for managing giant mimosa established.  Trials for managing giant mimosa established.  Invasive species monitoring and management plan developed and actively implemented by management board.  Management plan contribute to a part of the overall site management and zonation plan.  Mife protection and monitoring, with a focus on the Sarus Crane and zonation plan.  Zones in BPL area precisely delineated for different usage.  Establishment of a comanagement framework, with stronger involvement of local

	Increased patrol and enforcement efforts for wildlife protection and encroachment activities.				and community- focused NGOs)
Strengthen biodiversity monitoring, with a focus on the Sarus Crane population.	Increased awareness of Sarus Crane and wetland conservation amongst local people and tourists	100,000	3 years	Awareness activities on importance of preserving the cranes and the wetlands.  Biodiversity monitoring actively implemented.	
Component 3. Strengthen management of site fisheries, with a focus on small-scale artisanal fishers.	Fishing activities in BPL better regulated and enforced.  Management system strengthened, fishery resources are better managed and conserved.  Stronger management and protection of fish breeding areas.	500,000	5 years	Meetings and workshops to engage local stakeholders for participatory processes.  Licensing framework for fishers established and in place.  Patrol activities, using SMART approach against illegal fishing strengthened and scaled up significantly against baselines. Fishery monitoring framework in place.  Community led agreements on sustainable fishing practices,	Fisheries Administration (FiA)  BPL management board  Takeo provincial government  Borey Chulsar and Koh Andeth district governments  Fishing operators

				supported with improved storage and processing capability to increase market access and reduce wastage.			
Strengthen the capacity and resilience of small-scale artisanal fishers	Fishers (from the local communities of 12 villages) aware of wetland impacts, and adopt good practices.  Local fishing cooperatives and communities developed and have stronger access to livelihood opportunities.  Improved access to micro-finance for small scale fishing communities based around sustainable practices. Improved engagement of women in the workforce.	200,000	5 years	At least 100 small-scale fishing operators and fishers engaged across two districts.  Capacity building for local fishing communities (with a representative % of women) in cooperatives and sustainable fishery management.  Small loan scheme in place for small-scale fishers.  New fishing cooperatives established; existing cooperatives strengthened.	MAFF  Borey Chulsar and Koh Andeth district governments  Fishing operators  Community groups  Conservation organisations		
Component 4. Upscaling tourism infrastructure and strengthening sustainable ecotourism							
Improve and	Ecotourism	500,000	2 years	Ecotourism as a	MOE		
expand tourism	infrastructure		_	source of local			
infrastructure at	improved, to			livelihood	BPL		
Boeng Prek	enhance			strengthened,	management		
Lapouv,	opportunities			,	board		

including	for visitors to			benefiting at least	
signages and	observe Sarus			100 households.	Takeo
viewing	Cranes,				provincial
structures	Yellow-			Ecotourism strategy	government
	breasted			and business plans	
	Bunting and			and packages with	Borey Chulsar
	experience			a focus on Sarus	and Koh Andeth
	wetland			Crane and	district
	habitats			migratory	government
				waterbirds	
Strengthening	Improved	200,000	3 years	developed and	Conservation
the capacity of	capacity of			tested with tourism	and academic
local	local			operators.	organisations
communities and	stakeholders				(including bird
businesses for	to support			Stakeholder	and community-
nature-based	domestic and			consultation	focused NGOs)
tourism.	international			meetings	
	tourists.			organised.	Tourism
Establish a				Meetings and	operators
development				workshops to	
plan for nature-				engage local	
based tourism in				stakeholders for	
BPL				participatory	
				processes.	

Component 5. Strengthening sustainable and organic rice farming (on Jasmine rice, or other native rice varieties) in BPL

Piloting of	Successful	100,000	3 years	Successful rice-	Ministry of
sustainable,	trialing of pilot	100,000	o years	planting trials and	Agriculture
regenerative	plots of			experiments (on	Agriculture
rice-farming in	sustainable,			fertilizer use, soil	Takeo
Koh Andet and	regenerative			management) in	provincial
Borey Chulsar,	rice with			selected plots in	government
and in BPL	selected			BPL experimental	
experimental	communities in			·	Koh Andeth and
zone ('Ibis Rice'	BPL.			zone.	Borey Chulsar
and 'Crane Rice'	DFL.			Stakeholder	district
and Crane Rice				consultation	government
				meetings	Conservation
				organised.	organisations
				Meetings and	(including bird
				workshops to	and community-
				engage local	focused NGOs)
				stakeholders for	
				participatory	Agricultural
				processes.	banks
				Capacity building	
				(and training)	
				programmes for	
				organic and	
				biodiversity-friendly	
				rice farming.	
Scaling up of	Organic and	500,000	10 years	Stakeholder	Ministry of
sustainable,	biodiversity-			consultation	Agriculture
regenerative	friendly rice			meetings	Tales
rice-farming in	farming			organised.	Takeo provincial
the rice farming	expanded and			Meetings and	government
landscape that	upscaled in			workshops to	govorninon
surround BPL,	landscapes			engage local	Koh Andeth and
using organic	around BPL.			stakeholders for	Borey Chulsar
and/or wildlife-				participatory	district
friendly .	Financing			processes.	government
approaches.	mechanism				
	(for local loans			Exchange visits to	Representatives
	and grants)			Anlung Pring	from
	created for			wetlands to meet	households
	local			local communities	'Crane Rice'
	communities,			and 'Crane Rice'	community
	and supported			representatives.	groups
	by key lending			B	
	institutions for			Biodiversity-friendly	Conservation
	farmers.			rice-farming using	organisations
1	Incentives for			'Crane Rice' as a	

	biodiversity- friendly farming established.			source of local livelihood strengthened and increased by 50% of baselines. At least 30% of rice fields around BPL under organic/biodiversity-friendly cultivation.	Rural development organisations Fertilizer companies
Improved management of agro-chemical waste in the rice paddies in BPL experimental zone and surrounding areas in Koh Andeth and Borey Chulsar.	Reduced inflow of agrochemicals into the wetlands.  Increase in biodiversity in rice fields against baseline metrics.	500,000	3 years	Agro-chemical management and disposal protocols established and strengthened for all relevant villages.  Monitoring data shows reduced inflow of agro-chemicals into BPL wetlands wetlands against baselines.	

The proposed project is expected to be implemented over a period of at least five (5) years, with main project components focusing on improved site management for BPL (proposed to be led by the Ministry of Environment) and expanding organic and/or biodiversity-friendly rice agriculture (with the Ministry of Agriculture, Forestry and Fisheries). Conservation organisations are expected to play a major role in the project by supporting project activities focused on biodiversity monitoring, stakeholder engagement and capacity building for better management of fisheries and rice agriculture.

Assessment of impacts of project implementation has been proposed to assess how each component may benefit local communities or impact the project site (see Table 5). It is important to assess how major infrastructural developments in Takeo, such as canals and roads may potentially impact the site in the long-term.

The proposed interventions are expected to undertake activities promote gender inclusion and participation in livelihood activities, through capacity building activities for organic rice farming agriculture, small-scale fisheries and nature-based tourism.

There are no indigenous communities in the project landscape.

#### References

Asian Waterbird Census (AWC): Cambodia data for 2017-2022. See <a href="https://eaaflyway.net/asian-waterbird-census">https://eaaflyway.net/asian-waterbird-census</a>

BirdLife International Important Bird Area factsheet: Boeung Prek Lapouv (Cambodia). Available at: https://datazone.birdlife.org/site/factsheet/boeung-prek-lapouv-iba-cambodia

IUCN Red List of Threatened Species. Available at: www.iucnredlist.org

Ly S., Bunnath Z.S., Yong D.L., Bou V. & Chheoun S. (2022) Emerging evidence shows the global importance of the Boeung Prek Lapouv Protected Landscape, Cambodia for yellow-breasted buntings *Emberiza aureola*. *Cambodian Journal of Natural History* 2022: 78–83. Available at: <a href="https://www.fauna-flora.org/our-work/publications/cambodian-journal-natural-history/">https://www.fauna-flora.org/our-work/publications/cambodian-journal-natural-history/</a>

Mundkur, T. & Langendoen, T. (2022) Report on the Conservation Status of Migratory Waterbirds of the East Asian – Australasian Flyway Partnership. First Edition. Draft Report to the East Asian – Australasian Flyway Partnership. Wetlands International, Ede, The Netherlands. Available at: <a href="https://www.wetlands.org/publication/eaaf-conservation-status-review1/">https://www.wetlands.org/publication/eaaf-conservation-status-review1/</a>

NatureLife Cambodia / BirdLife International Cambodia Programme (2022) *Annual report 2022: highlight key achievements*. Available at: <a href="https://naturelifecambodia.org/wp-content/uploads/2023/06/NLC-2022-annual-report-Final-Email.pdf">https://naturelifecambodia.org/wp-content/uploads/2023/06/NLC-2022-annual-report-Final-Email.pdf</a>

Pinn, T., Chhoeum, C., & Saman, N. (2012). An Aassessment of the Natural Resources of Ang Trapeang Thmor and Its Impact on the Livelihoods of Pongro and Sambour Villages. *International Journal of Environmental and Rural Development*, *3*(2), 72-76.

Seng Kim Hout, Pech Bunnat, Poole, C.M., Tordoff, A.W., Davidson, P. and E. Delattre (2003) *Directory of Important Bird Areas in Cambodia: key sites for conservation.* Phnom Penh, Cambodia: Department of Forestry and Wildlife, Department of Nature Conservation and Protection, BirdLife International in Indochina, and the Wildlife Conservation Society Cambodia Program.

Sophanna, L. (2017). Project report 2014-2017. Wildfowl & Wetlands Trust, Cambodia.

Sophanna, L., Pok, H. and Avent, T. (2019) Climate change vulnerability assessment for Boueng Prek Lapouv Protected Landscape, Cambodia. Bangkok, Thailand: IUCN ARO. X + 32pp. Available at:

https://iucn.org/sites/default/files/content/documents/2019/climate\_change\_vulnerability\_assess ment\_boeung\_prek\_lapouv\_protected\_landscape\_cambodia.pdf

van Zalinge, R., Ses Vong Sambat, Peh, K., Um Sokun, Graham, A., & Seng Kim Hout (2013) *Consolidated results of an ecosystem service assessment for Boeung Prek Lapouv*. Wildfowl & Wetlands Trust (WWT), Cambodian Lower Mekong Wetlands Project, Phnom Penh.

WWT (2014). Boeung Prek Lapouv Management Plan: January 2014 – December 2018. Wildfowl & Wetlands Trust, Cambodia. Available at: <a href="https://www.cepf.net/sites/default/files/bpl">https://www.cepf.net/sites/default/files/bpl</a> mp 2013 2018 240613 1.pdf

#### **Annex 1**. Supplementary information on flood mitigation services

To further validate the identification of the top ecosystem services by means of stakeholder consultation, an expectedly essential or non-substitutable regulating service across all RFI sites, namely coastal protection and flood mitigation (i.e., storm and flood hazard regulation), was assessed based on a combination of globally available datasets supplemented by web-based tool Co\$tingNature (Mulligan, 2022). Estimates for flood mitigation were spatially inferred in QGIS from a selection of metrics expressing different biophysical values modelled online by the Water World component of this tool. Equivalent data to assess monetary values similarly to coastal protection were not available for the RFI region.

The key metrics selected for biophysical values (Table A1) were the average green storage capacity, which is the volume of water stored by each square kilometre of wetland itself as well as its soil and vegetation, and the direct influence of this storage capacity on beneficiaries found downstream of the wetland, both as the average number of people and the average built-up area that are uniquely benefitting from the resulting flood mitigation (and not from other green storage found upstream).

**Table A1.** Contribution of the wetland habitats to flood mitigation in Boeung Prek Lapouv based on site-level (biophysical) values inferred from Mulligan (2022) and expressed as ranges to represent the resulting uncertainty.

Influence of the wetland on flood mitigation (metrics)	Benefit/Beneficiaries
Average green storage capacity per sq. km of wetland in million cubic metres (GigaLitres/km²)	410 – 485
Average population uniquely benefitting from influential green storage upstream per sq. km of wetland (n/km²)	127 – 150
Average built-up area uniquely benefitting from influential green storage upstream per sq. km of wetland (ha/km²)	7.69 – 9.11

**Table A2.** Biophysical benefits from and beneficiaries of RFI inland wetland sites (expressed as ranges to represent the resulting uncertainty) and at the national level.

Site name	Green storage capacity (GigaLitres/km²)	Downstream population (n /km²)	Downstream built-up area (ha /km²)	
Prek Toal Core Area	486 (±29)	112 (±7)	7.19 (±0.42)	
Ang Tropeang Thmor	411 (±38)	73 (±7)	3.84 (±0.36)	
Boeung Prek Lapouv	448 (±37)	139 (±12)	8.40 (±0.71)	
Anlung Pring	264 (±63)	0 (±0)	1.22 (±0.29)	
Boeng Tonle Chhmar	406 (±27)	102 (±7)	6.57 (±0.43)	

Chikraeng and Stoung	206 (±23)	45 (±5)	2.84 (±0.32)	
Kulen Promtep Wildlife	456 (±6)	87 (±1)	4.20 (±0.05)	
Sanctuary				
Cambodia RFI average	384	81	4.89	
Cambodia national average	399	109	5.24	

**Table A5**. Key habitat types in Boeung Prek Lapouv (Sarus Crane Reserve) based on stakeholder-based assessment at the Regional Flyway Initiative workshop in September 2023.

	• • •		•		
Lightet type	Current	state	Alternative state (2035)		
Habitat type	Area (ha)	Cover (%)	Area (ha)	Cover (%)	
Permanent freshwater lakes	305.2	3.7	105.1	1.3	
Planted/Cultivated (non-woody crops)/Others	6004.9	72.2	5924.8	71.3	
Grasslands	1501.2	18.1	1501.2	18.1	
Freshwater, tree-dominated wetlands	500.4	6.0	700.6	8.4	
Low Intensity Residential	0.0	0.0	80.1	1.0	
Total	8,311.8	100.0	8,311.8	100.0	